

PE- CONJUGATED MONOCLONAL ANTIBODY TO MOUSE MONOCYTE, MACROPHAGE MARKER (CD14)

clone Sa14-2



Catalog no HM1060PE-25 (lot number and expiry date are indicated on the label)

Description The monoclonal antibody Sa14-2 recognizes the mouse monocyte marker CD14. The CD14 receptor is a pattern recognition molecule in the innate immune response against microorganisms and other exogenous and endogenous stress factors. CD14 was characterized as a receptor for LPS. The CD14 gene consists of two exons which code for a single mRNA that is translated into a protein of 375 amino acids. The CD14 protein is composed of eleven leucin-rich repeats, which are also found in TLR and which are important in PAMP binding. In contrast to TLR, however, CD14 lacks a transmembrane domain, and thus cannot initiate intracellular signal transduction by itself. The most important CD14 signaling co-receptor is toll-like receptor 4 (TLR4), which activates, among others, the nuclear factor κB (NF-κB) inflammatory pathway. The CD14 protein is processed in the endoplasmatic reticulum and expressed as a 55 kDa glycoprotein on the cell surface via a glycosylphosphatidyl (GPI) anchor. Like other GPI-anchored proteins, CD14 accumulates on the cell surface in microdomains known as lipid rafts. CD14 is expressed pre dominantly on the surface of 'myeloid' cells, such as monocytes, macrophages and neutrophils, but at lower levels also on epithelial cells, endothelial cells and fibroblasts. CD14 is also expressed in a soluble form (sCD14). sCD14 may result from secretion of the protein before coupling to the GPI anchor or from shedding or cleavage from the surface of monocytes. sCD14 is present in the circulation and other body fluids and levels of sCD14 in plasma increase during inflammation and infection. CD14 is a molecule with a broad range of functions. In addition to functioning as a pattern recognition receptor for a variety of microbial ligands, CD14 also acts as a receptor for endogenous molecules like intercellular adhesion molecule (ICAM)-3 on the surface of apoptotic cells, amyloid peptid, ceramide, and urate crystals. Ligation of CD14 by these ligands, except for apoptotic cells, mediates activation of inflammatory responses.

Aliases Monocyte differentiation antigen CD14

Immunogen CD14 transfectant

Species Rat IgG2a


Formulation 25 tests 0.2 µm filtered antibody solution in PBS, containing 1% bovine serum albumin and 0.02% sodium azide.

Application

	F	FC ³	FS	IA	IF ⁴	IP	P	W
Yes		•	•		•			
No								
N.D.	•			•		•	•	•

N.D.= Not Determined; *F* = Frozen sections; *FC* = Flow Cytometry; *FS* = Functional Studies; *IA* = Immuno Assays; *IF* = Immuno Fluorescence; *IP* = Immuno Precipitation; *P* = Paraffin sections; *W* = Western blot

Application notes



References	<ol style="list-style-type: none"> 1. Akashi, S et al; Lipopolysaccharide interaction with cell surface Toll-like receptor 4-MD-2: higher affinity than that with MD-2 or CD14. <i>J. Exp. Med.</i> 2003, 198:1035-1042 2. Hajshengallis, G et al; Porphyromonas gingivalis fimbriae proactively modulate beta2 integrin adhesive activity and promote binding to and internalization by macrophages. <i>Infect Immun</i> 2006, 74: 5658 3. Kiyokawa, T et al. A single base mutation in the PRAT4A gene reveals differential interaction of PRAT4A with Toll-like receptors. <i>Int Immunol</i> 2008, 20:1407-1415 4. Wilkinson, T et al; Trappin-2 promotes early clearance of pseudomonas aeruginosa through CD14-dependent macrophage activation and neutrophil recruitment. <i>Am J Pathol</i> 2009, 174: 1338 5. Bi, S et al. Galectin-9 binding to cell surface protein disulfide isomerase regulates the redox environment to enhance T-cell migration and HIV entry. <i>PNAS</i> 2011, 108:10650 										
Use	For flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The recommended use of this reagent is 10 µl per 250.000 cells in a 100 µl total staining volume.										
Positive control	RAW-cells										
Storage and stability	Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year. The exact expiry date is indicated on the label.										
Precautions	For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent infringements that might result from the use or derivation of this product.										
Also available	<table border="0" style="width: 100%;"> <tr> <td style="width: 20%;">HM1060</td> <td>Monoclonal antibody against Mouse CD14, clone Sa14-2</td> </tr> <tr> <td>HM1060F</td> <td>FITC conjugated monoclonal antibody against Mouse CD14, clone Sa14-2</td> </tr> <tr> <td>HM1060PE100</td> <td>PE conjugated monoclonal antibody against Mouse CD14, clone Sa14-2</td> </tr> <tr> <td>HM1029</td> <td>Monoclonal antibody against mouse TLR4/MD2, clone MTS510</td> </tr> <tr> <td>HM2060</td> <td>Monoclonal antibody against human CD14, clone MEM-15</td> </tr> </table>	HM1060	Monoclonal antibody against Mouse CD14, clone Sa14-2	HM1060F	FITC conjugated monoclonal antibody against Mouse CD14, clone Sa14-2	HM1060PE100	PE conjugated monoclonal antibody against Mouse CD14, clone Sa14-2	HM1029	Monoclonal antibody against mouse TLR4/MD2, clone MTS510	HM2060	Monoclonal antibody against human CD14, clone MEM-15
HM1060	Monoclonal antibody against Mouse CD14, clone Sa14-2										
HM1060F	FITC conjugated monoclonal antibody against Mouse CD14, clone Sa14-2										
HM1060PE100	PE conjugated monoclonal antibody against Mouse CD14, clone Sa14-2										
HM1029	Monoclonal antibody against mouse TLR4/MD2, clone MTS510										
HM2060	Monoclonal antibody against human CD14, clone MEM-15										