

**CERTIFICATE OF ANALYSIS – TECHNICAL DATA SHEET**

<b>Product name</b>	Ethenoadenosine, clone 1G4		
<b>Catalog number</b>	HM5005		
<b>Lot number</b>	-	<b>Expiry date</b>	-
<b>Volume</b>	1 ml	<b>Amount</b>	100 µg
<b>Formulation</b>	0.2 µm filtered in PBS+0.1%BSA+0.02%NaN3	<b>Concentration</b>	100 µg/ml
<b>Host Species</b>	Mouse IgG2	<b>Conjugate</b>	None
<b>Endotoxin</b>	N.A.	<b>Purification</b>	Protein G
<b>Storage</b>	4°C		

**Application notes**

	IHC-F	IHC-P	IF	FC	FS	IA	IP	W
Reference #								
Yes	•			•		•	•	•
No								
N.D.		•	•		•			

*N.D.= Not Determined; IHC = Immuno histochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot*

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:10.

**General Information**

<b>Description</b>	The reaction of chloroacetaldehyde (CAA) with DNA results in the formation of ethenoadenosine also known as 1,N <sup>6</sup> -ethenodeoxy-adenosine or etheno-A. CAA is formed through metabolization of for example vinyl chloride, a well-established carcinogen. In RNA, after in vitro activation, etheno-A and etheno-C are the principle products of RNA damage. Etheno derivate formation may be highest in single-stranded DNA-regions. The monoclonal antibody 1G4 reacts with both the ribose and deoxyribose form of the adduct. The antibody is not cross reactive with non-modified DNA or the normal nucleotides.
<b>References</b>	<ol style="list-style-type: none"> <li>Young, T et al; Development of techniques to monitor for exposure to vinyl chloride: monoclonal antibodies to ethenoadenosine and ethenocytidine. Carcinogenesis 1988, 9: 589</li> <li>Krebs, C et al; Flow cytometric and immunoblot assays for cell surface ADP-ribosylation using a monoclonal antibody specific for ethenoadenosine. Anal Biochem 2003, 314: 108</li> </ol>
<b>Storage&amp;stability</b>	Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year.
<b>Precautions</b>	For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent infringements that might result from the use or derivation of this product.

We hereby certify that the above-stated information is correct and that this product has been successfully tested by the Quality Control Department. This product was released for sale according to the existing specifications. This document has been produced electronically and is valid without a signature.

Approved by Manager of QC  
Robbert Zwinkels

Date  
16/03/2018

Do you have any questions or comments regarding this product? Please contact us via [support@hycultbiotech.com](mailto:support@hycultbiotech.com).