

CERTIFICATE OF ANALYSIS – TECHNICAL DATA SHEET

Product name TNF-RII, Mouse, pAb

Catalog number	HP8003		
Lot number	-	Expiry date	-
Volume	1 ml	Amount	100 μg
Formulation	0.2 μ m filtered in PBS+0.1%BSA	Concentration	100 μg/ml
Host species	Rabbit IgG	Conjugate	None
Endotoxin	<24 EU/mg	Purification	Protein A
Storage	4°C		

Application notes

		IHC-F	IHC-P	IF	FC	FS	IA	IP	W
	Reference #								
	Yes				•	•	•	•	•
	No								
-	N.D.	•	•	٠					

N.D.= Not Determined; IHC = Immuno histochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.

FS: For functional studies, *in vitro* dilutions have to be optimized in user's experimental setting. HP8003 is cross reactive with Mouse TNF-RI which is problematic in case of TNF-RII knockout studies and when used in high concentrations in functional studies.

General Information

Description	The polyclonal antibody recognizes the extracellular part of the mouse Tumor Necrosis Factor Receptor type 2 (TNF- RII) of the membrane-bound as well as the soluble receptor. TNF-RII (~75-80 kDa) is present on most cell types and is considered to play a prominent role in cell stimulation by TNF-alpha. TNF-alpha activates inflammatory responses, induces apoptosis, regulates cellular proliferation, and may even promote cancer progression. The effects of TNF- alpha are mediated by TNF-RI and TNF-RII, which have both distinct and overlapping downstream signaling cascades. Induction of cytotoxicity and other functions are mediated largely via TNF-RI. TNF-RI is equally well activated by both the 17 kDa soluble and 26 kDa membrane-bound form, whereas TNF-RII is efficiently activated only by the membrane bound form of TNF-alpha. Binding of the inherently trimeric TNF-alpha to TNFR1 and TNFR2 induces receptor trimerization and recruitment of several signaling proteins to the cytoplasmic domains of the receptors. Occupancy of TNFR2 results in direct recruitment of TNF Receptor Associated Factor 2 (TRAF2), which in turn recruits TRAF1.			
Aliases	Tumor necrosis factor receptor superfamily member 1B, TNF-RII, p75/p80, CD120b			
Cross reactivity	Mouse TNF-RI: Yes; Mouse TNF-alpha: minimal			
References	 Lucas, R et al; Both TNF receptors are required for direct TNF-mediated cytotoxicity in microvascular endothelial cells. Eur J Immunol 1998, <i>28</i>: 3577 Bemelmans, M et al; LPS-Induced sTNF-receptor release in vivo in a murine model. J Immunol 1993, <i>151</i>: 5554 Gerspach, J et al; Detection of membrane-bound tumor necrosis factor (TNF): an analysis of TNF-specific reagents. Micr Res Tech 2000, <i>50</i>: 243 			

Storage&stability Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year.

Precautions For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent infringements that might result from the use or derivation of this product.

We hereby certify that the above-stated information is correct and that this product has been successfully tested by the Quality Control Department. This product was released for sale according to the existing specifications. This document has been produced electronically and is valid without a signature.

Approved by Manager of QC Robbert Zwinkels

Date 13/03/2018

Do you have any questions or comments regarding this product? Please contact us via support@hycultbiotech.com.

bringing innate immunity to the **next level**