


CERTIFICATE OF ANALYSIS – TECHNICAL DATA SHEET

| | | | |
|-----------------------|--|----------------------|-----------|
| Product name | I-FABP, Human, pAb | | |
| Catalog number | HP9020 | | |
| Lot number | - | Expiry date | - |
| Volume | 1 ml | Amount | 100 µg |
| Formulation | 0.2 µm filtered in PBS+0.1%BSA+0.02%NaN3 | Concentration | 100 µg/ml |
| Host Species | Rabbit IgG | Conjugate | None |
| Endotoxin | N.A. | Purification | Protein G |
| Storage | 4°C | | |

Application notes

| | IHC-F | IHC-P | IF | FC | FS | IA | IP | W |
|-------------|-------|-------|----|----|----|----|----|---|
| Reference # | | | | | | | | |
| Yes | | • | | | | • | | • |
| No | | | | | | | | |
| N.D. | • | | • | • | • | | • | |

N.D.= Not Determined; IHC = Immunohistochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot



IHC-P: Immunohistochemical staining of human jejunum paraffin embedded slides.

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.

General Information

| | |
|------------------------------|---|
| Description | The polyclonal antibody recognizes human intestinal fatty acid binding protein (I-FABP) of both natural and recombinant origin. The I-FABP protein is derived from the human FABP2 gene. FABPs are small intracellular proteins (~13-14 kDa) with a high degree of tissue specificity that bind long chain fatty acids. They are abundantly present in various cell types and play an important role in the intracellular utilization of fatty acids, transport and metabolism. There are at least nine distinct types of FABP, each showing a specific pattern of tissue expression. Due to its small size, FABP leaks rapidly out of ischemically damaged necrotic cells leading to a rise in serum levels. Ischemically damaged tissues are characterized histologically by absence (or low presence) of FABP facilitating recognition of such areas. I-FABP is localized in the small bowel epithelium, with highest expression level in the jejunum. |
| Aliases | FABP2, FABPI, Intestinal Fatty Acid Binding Protein |
| Cross reactivity | Rat I-FABP: Weak; Mouse I-FABP: Weak |
| Storage&stability | Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year. |

Precautions

For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent infringements that might result from the use or derivation of this product.

We hereby certify that the above-stated information is correct and that this product has been successfully tested by the Quality Control Department. This product was released for sale according to the existing specifications. This document has been produced electronically and is valid without a signature.

Approved by Manager of QC
Robbert Zwinkels

Date
13/03/2018

Do you have any questions or comments regarding this product? Please contact us via support@hycultbiotech.com.