


CERTIFICATE OF ANALYSIS – TECHNICAL DATA SHEET

Product name	C3/C3a des arg, Human, clone 2991					
Catalog number	HM2074-20UG					
Lot number	xxxxXXxxxx	Expiry date	MMM YYYY			
Volume	200 μΙ	Amount	20 µg			
Formulation	0.2 μm filtered in PBS+0.1%BSA+0.02%NaN3	Concentration	100 μg/ml			
Host Species	Mouse IgG1	Conjugate	None			
Endotoxin	N.A.	Purification	Protein G			
Storage	4°C					

Application notes

	IHC-F	IHC-P	IF	FC	FS	IA	IP	W
Reference #								
Yes						•		•
No								
N.D.	•	•	•	•	•		•	

N.D.= Not Determined; IHC = Immuno histochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot



IA: Three different batches tested in a sandwich ELISA to determine the specificity of HM2074. HM2074 was used as detection antibody.

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.

- W: A reduced sample treatment and SDS-PAGE was used. The band size is ~8 kDa.
- Positive control: Activated human serum or purified human C3a desArg protein; Negative control: For Western blot, any irrelevant protein can be used.

General Information

Description The monoclonal antibody 2991 recognizes the neo-epitope of human C3a/C3a-desArg that is formed upon cleavage of the complement protein C3 into C3a and C3b. The complement system is an important part of the humoral response in innate immunity, consisting of three different pathways. The third complement component, C3, is central to the classical, alternative and lectin pathways of complement activation. Activation products of the complement cascade contain neo-epitopes that are not present in the individual native components. Monoclonal antibodies detecting neo-epitopes have been used for direct quantification of activation at different steps in the complement cascade. The synthesis of C3 is tissue-specific and is modulated in response to a variety of stimulatory agents. C3 is the most abundant protein of the complement system with serum protein levels of about 1.3 mg/ml. An inherited deficiency of C3 predisposes the person to frequent assaults of bacterial infections. In ulcerative colitis, and idiopathic chronic inflammatory bowel disease, the deposition of C3 in the diseased mucosa has been reported. Proteolysis by certain enzymes results in the cleavage of C3 releasing C3a anaphylatoxin and C3b. C3a is a protein of 74 amino acids. C3a itself is very short-lived and in serum is cleaved rapidly into the more stable C3a-desArg (also called acylation stimulating protein, ASP). Therefore, measurement of C3a-desArg allows reliable conclusions about the level of complement activation in the test samples. C3a is a mediator of local inflammatory processes. It induces smooth



muscle contraction, increases vascular permeability, and causes histamine release from mast cells and basophilic leukocytes. C3a is involved in inflammatory reactions seen in gram-negative bacterial sepsis, trauma, ischemic heart disease, post-dialysis syndrome and a variety of autoimmune diseases. Normal values in plasma of control persons ranges between 48 – 150 ng/ml (median: 86.4 ng/ml, SD: 29.3 ng/ml). Monoclonal antibody 2991 recognizes both C3a and C3a-desArg, with at least a 5x higher preference for C3a-desArg.

Immunogen	Human C3a desArg
Aliases	Acylation stimulating protein, ASP
Gene	Gene name: C3
Storage&stability	Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year.
Precautions	For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent infringements that might result from the use or derivation of this product.

We hereby certify that the above-stated information is correct and that this product has been successfully tested by the Quality Control Department. This product was released for sale according to the existing specifications. This document has been produced electronically and is valid without a signature.

Approved by Manager of QC

Date

Do you have any questions or comments regarding this product? Please contact us via support@hycultbiotech.com.