


CERTIFICATE OF ANALYSIS – TECHNICAL DATA SHEET

| | | | |
|-----------------------|--|----------------------|-----------|
| Product name | PECAM-1, Mouse, clone ER-MP12 | | |
| Catalog number | HM1084-200UG | | |
| Lot number | - | Expiry date | - |
| Volume | 200 µl | Amount | 20 µg |
| Formulation | 0.2 µm filtered in PBS+0.1%BSA+0.02%NaN3 | Concentration | 100 µg/ml |
| Host Species | Rat IgG2a | Conjugate | None |
| Endotoxin | N.A. | Purification | Protein G |
| Storage | 4°C | | |

Application notes

| | IHC-F | IHC-P | IF | FC | FS | IA | IP | W |
|-------------|-------|-------|----|---------|----|----|-----|---|
| Reference # | 3 | | | 1,2,4-7 | | 1 | 1,4 | |
| Yes | • | • | | • | | • | • | |
| No | | | | | | | | |
| N.D. | | | • | | • | | | • |

N.D.= Not Determined; IHC = Immuno histochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot



IHC-F: Mouse spleen frozen sections were stained with ER-mp12 (2µg/ml).

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.

- Positive control: Macrophage cell line RMB-1.

General Information
Description

The monoclonal antibody ER-MP12 recognizes mouse platelet endothelial cell adhesion molecule-1 (PECAM-1;CD31). PECAM-1 is a 130kDa glycoprotein involved in cell adhesion and a signalling receptor that is expressed on hematopoietic and endothelial cells. It is a member of the immunoglobulin (Ig)-superfamily of cell adhesion molecules. The type I transmembrane protein is composed of an extracellular region of six Ig-like homology domains, a 19-residue transmembrane domain, and a 118 residue cytoplasmic tail and is differentially glycosylated involving N-linked and O-linked glycosylation sites. The signalling of PECAM-1 is based on 2 immuno receptor tyrosine inhibitory motifs (ITIMs) in its cytoplasmic domain. These ITIM domains can serve as docking sites for signalling molecules such as protein tyrosine phosphatases, and ligation of PECAM-1 can induce phosphorylation of tyrosine- and serine/threonine-residues of these intracellular regions eventually leading to activation of signalling pathways. The biological characteristics of PECAM-1 in these pathways and cellular adhesion have been mapped to specific regions of the PECAM-1 molecule.

PECAM-1 is abundantly expressed on endothelial cells, where it is a major constituent of the endothelial cell intercellular junction in confluent vascular beds, making it a commonly used endothelial marker. Furthermore, it is expressed on most cells of the hematopoietic lineage including platelets, monocytes, neutrophils, and lymphocyte subsets. Besides adhesive properties PECAM-1 is an efficient signalling molecule. The protein is involved in vascular biology including angiogenesis, platelet function, and thrombosis, mechanosensing of endothelial cells, and regulation of multiple stages of leukocyte migration through venular walls. PECAM-1 is also a regulator of inflammatory responses. It has been shown to serve a variety of pro-inflammatory and anti-inflammatory functions. Pro-inflammatory functions of PECAM-1 include the facilitation of leukocyte transendothelial migration and the transduction of mechanical signals in endothelial cells originating from fluid shear stress. Anti-inflammatory functions include the

dampening of leukocyte activation, suppression of pro-inflammatory cytokine production, and the maintenance of vascular barrier integrity. Monoclonal antibody ER-MP12 reacts also with the major population of colony forming unit macrophage (CFU-M) precursor cells, with a subpopulation of pre CFU-M and monoblasts, and with precursor cells of granulocytes. The antibody can also be used, in combination with the monoclonal antibody ER-MP20, for identification of different stages of myeloid progenitor cell development.

| | |
|------------------------------|--|
| Immunogen | Mouse macrophage precursor hybrid cells. W1C3 (Ref.1) |
| Aliases | Platelet endothelial cell adhesion molecule-1; CD31; EndoCam |
| References | <ol style="list-style-type: none">1. Leenen, P et al; Murine macrophage precursor characterization II. Monoclonal antibodies against macrophage precursor antigens. <i>Eur J Immunol</i> 1990, <i>20</i>: 272. De Bruijn, M et al; Distinct mouse bone marrow macrophage precursors identified by differential expression of ER-MP12 and ER-MP20 antigens. <i>Eur J Immunol</i> 1994, <i>24</i>: 22793. Morioka, Y et al; Immunophenotypic and ultrastructural heterogeneity of macrophage differentiation in bone marrow and fetal hematopoiesis of mouse in vitro and in vivo. <i>J Leukoc Biol</i> 1994, <i>55</i>: 6424. Ling, V et al; Structural identification of the hematopoietic progenitor antigen ER-MP12 as the vascular endothelial adhesion molecule PECAM-1 (CD31). <i>Eur J Immunol</i> 1997, <i>27</i>: 5095. De Bruijn, M et al; Bone marrow cellular composition in <i>Listeria monocytogenes</i> infected mice detected using ER-MP12 and ER-MP20 antibodies: a flow cytometric alternative to differential counting. <i>J Immunol Methods</i> 1998, <i>217</i>: 276. Van Rijt, L et al. Allergen-induced accumulation of airway dendritic cells is supported by an increase in CD31hiLy-6Cneg bone marrow precursors in a mouse model of asthma. <i>Blood</i> 2002, <i>100</i> :36637. Jin, M et al. Enhancement of repopulation and hematopoiesis of bone marrow cells in irradiated mice by oral administration of PG101, a water-soluble extract from <i>Lentinus lepideus</i>. <i>Exp biol med</i> 2003, <i>228</i> :759 |
| Storage&stability | Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year. |
| Precautions | For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent infringements that might result from the use or derivation of this product. |

We hereby certify that the above-stated information is correct and that this product has been successfully tested by the Quality Control Department. This product was released for sale according to the existing specifications. This document has been produced electronically and is valid without a signature.

Approved by Manager of QC
Brenda Teunissen

Date
28/10/2020

Do you have any questions or comments regarding this product? Please contact us via support@hycultbiotech.com.