

## **CERTIFICATE OF ANALYSIS – TECHNICAL DATA SHEET**

Product name	Elafin/SKALP, Human, clone TRABO2					
Catalog number	HM2062-100UG					
Lot number	-	Expiry date	-			
Volume	1 ml	Amount	100 μg			
Formulation	0.2 $\mu m$ filtered in PBS+0.1%BSA+0.02%NaN3	Concentration	100 μg/ml			
Host Species	Mouse IgG1	Conjugate	None			
Endotoxin	N.A.	Purification	Protein G			
Storage	4°C					

## **Application notes**

	IHC-F	IHC-P	IF	FC	FS	IA	IP	W
Reference #								
Yes		•				•		•
No	•							
N.D.			•	•	•		•	

N.D.= Not Determined; IHC = Immuno histochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:10.

W: For Western blotting it is recommended to use non-reducing conditions.

## **General Information**

Description	lafin is an epithelial proteinase inhibitor also known under various other names such as Skin-derived Anti pukoproteinase (SKALP) and Elastase-Specific Inhibitor (ESI). Elafin belongs to the Trappin gene family and was iven the systematic name Trappin-2. The Trappin family is defined by a N-terminal transglutaminase substrate domain nd a C-terminal four disulphide core. Trappins have been suggested to play a role in the regulation of inflammation nd in protection against tissue damage in stratified epithelia. Elafin is an inhibitor of leukocyte elastase and proteinase- and is a substrate for transglutaminases. The protein is constitutively expressed in various epithelia including hair plices, oesophagus, vagina and oral cavity. Elafin is not present in normal human skin but is strongly induced during flammation as in psoriasis and wound healing. Antibodies to Elafin can be used to evaluate the effects of treatment f psoriasis since its expression is significantly correlated with clinical scores. Antibodies to Elafin have also been uccesfully used to study differentiation in squamous cell carcinoma of the head-and-neck region, oesophagus and kin. It is also shown that Elafin possesses antimicrobial activity against gram-positive and gram-negative bacteria. RAB2O recognises all known Trappins via the GQDPVK epitope. TRAB2O shows strong reaction with human native ull-length Elafin/SKALP and a weaker reaction with fully processed Elafin/SKALP (C-terminal 57 aa). For detection of ulty processed Elafin/SKALP (C-terminal 57 aa) we advise to use HM2063. The antibody is cross reactive with Trappin amily members of other species with GQDPVK epitope.			
Cross reactivity	Other species with GQDPVK epitope: Yes.			
References	<ol> <li>Vandermeeren, M et al; Development and application of monoclonal antibodies against SKALP/elafin and other trappin family members. Arch Dermatol Res 2001, <i>293</i>: 343</li> <li>Alkemade, J et al; Demonstration of skin-derived antileukoproteinase (SKALP) and its target enzyme human leukocyte elastase in squamous carcinoma. J Pathol 1994, <i>174</i>: 121</li> <li>Bergen van, H et al; Expression of SKALP/elafin during wound healing in human skin. Arch Dermatol Res 1996, <i>288</i>: 458</li> <li>Pfundt, R et al; Constitutive and inducible expression of SKALP/elafin provides anti-elastase defense in human epithelia. J Clin Invest 1996, <i>98</i>: 1389</li> <li>Schalkwijk, J et al; Localization of SKALP/elafin. J Invest Dermatol 1993, <i>100</i>: 390</li> <li>Schalkwijk, J et al; The trappin gene family: proteins defined by an N-terminal transglutaminase substrate domain and a c-terminal four-disulphide core. Biochem J 1999, <i>340</i>: 569</li> <li>Wiedow, O et al; Elafin: an elastase-specific inhibitor of human skin. J Biol Chem 1990, <i>265</i>: 14791</li> <li>Huh, WK et al; Dynamic alteration of human beta-defensin 2 localization from cytoplasm to intercellular space in psoriatic skin. J Mol Med 2002, <i>80</i>: 678</li> </ol>			
Storage&stability	Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year.			

## Precautions

For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent infringements that might result from the use or derivation of this product.

We hereby certify that the above-stated information is correct and that this product has been successfully tested by the Quality Control Department. This product was released for sale according to the existing specifications. This document has been produced electronically and is valid without a signature.

Approved by Manager of QC Brenda Teunissen

Date 29/11/2019

Do you have any questions or comments regarding this product? Please contact us via support@hycultbiotech.com.