

## **CERTIFICATE OF ANALYSIS – TECHNICAL DATA SHEET**

| Product name   | C3/C3a/C3a-desArg, Human, clone 2898          |               |           |  |  |  |  |
|----------------|---|---------------|-----------|--|--|--|--|
| Catalog number | HM2075-20UG                                   |               |           |  |  |  |  |
| Lot number     | -   | Expiry date   | -         |  |  |  |  |
| Volume         | 200 μΙ  | Amount        | 20 µg     |  |  |  |  |
| Formulation    | $0.2~\mu m$ filtered in PBS+0.1%BSA+0.02%NaN3 | Concentration | 100 μg/ml |  |  |  |  |
| Host Species   | Mouse IgG1                                    | Conjugate     | None      |  |  |  |  |
| Endotoxin      | N.A.  | Purification  | Protein G |  |  |  |  |
| Storage        | 4°C   |               |           |  |  |  |  |

## **Application notes**

|             | IHC-F | IHC-P | IF | FC | FS | IA | IP | W   |
|-------------|-------|-------|----|----|----|----|----|-----|
| Reference # |       |       |    |    |    |    |    | 1,2 |
| Yes         |       |       |    |    |    | •  |    | •   |
| No          |       |       |    |    |    |    |    |     |
| N.D.        | •     | •     | •  | •  | •  |    | •  |     |

N.D.= Not Determined; IHC = Immuno histochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.

- W: A reduced sample treatment and SDS-PAGE was used. The band size is ~95 and 8.2 kDa for C3 and C3a respectively (Ref.1).
- Positive control: Human plasma; Negative control: C3-depleted serum.

## **General Information**

| Description       | The monoclonal antibody 2898 recognizes the C-terminus of human complement protein C3a. The antibody recognize intact C3, C3a as well as C3a-desArg. C3 is the most abundant protein of the complement system with serum protei levels of about 1.3 mg/ml. The complement system is an important part of the humoral response in innate immunity consisting of three different pathways. The third complement component, C3, is central to the classical, alternative an lectin pathways of complement activation. Activation products of the complement cascade contain neo-epitopes that are not present in the individual native components. Monoclonal antibodies detecting neo-epitopes have been used for direct quantification of activation at different steps in the complement cascade. The synthesis of C3 is tissue-specific and is modulated in response to a variety of stimulatory agents. An inherited deficiency of C3 predisposes the perso to frequent assaults of bacterial infections. In ulcerative colitis, and idiopathic chronic inflammatory bowel disease, th deposition of C3 in the diseased mucosa has been reported. Proteolysis by certain enzymes results in the cleavage of C3 releasing C3a anaphylatoxin and C3b. C3a is a protein of 74 amino acids. C3a itself is very short-lived and in serur is cleaved rapidly into the more stable C3a-desArg (also called acylation stimulating protein, ASP). Therefore, measurement of L03a inflammatory processes. It induces smooth muscle contraction, increases vascula permeability, and causes histamine release from mast cells and basophilic leukocytes. C3a is involved in inflammatory reactions seen in gram-negative bacterial sepsis, trauma, ischemic heart disease, post-dialysis syndrome and a variet of autoimmune diseases. |  |  |  |  |
|-------------------|---|--|--|--|--|
| Immunogen         | Peptide CARASHLGLA (last nine amino acids of C3a-desArg)  |  |  |  |  |
| Aliases           | Complement protein C3   |  |  |  |  |
| References        | <ol> <li>Hartmann, H et al; Rapid quantification of C3a and C5a using a combination of chromatographic and<br/>immunoassay procedures. J Immunol Meth 1993, <i>166</i>: 35</li> <li>Smalley, D et al; Proteomic discovery of 21 proteins expressed in human plasma-derived but not platelet-derived<br/>microparticles. Thromb Haemost 2007, <i>97</i>: 67</li> </ol>   |  |  |  |  |
| Storage&stability | Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year.   |  |  |  |  |

## Precautions

For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent infringements that might result from the use or derivation of this product.

We hereby certify that the above-stated information is correct and that this product has been successfully tested by the Quality Control Department. This product was released for sale according to the existing specifications. This document has been produced electronically and is valid without a signature.

Approved by Manager of QC Brenda Teunissen

Date 16/11/2020

Do you have any questions or comments regarding this product? Please contact us via support@hycultbiotech.com.

bringing innate immunity to the next level