

CERTIFICATE OF ANALYSIS - TECHNICAL DATA SHEET

Product name C5aR, Human, clone W17/1

Catalog number HM2095-20UG

Lot number - Expiry date -

Volume 200 μl Amount 20 μg

Formulation 0.2 μm filtered in PBS+0.1%BSA+0.02%NaN3 Concentration 100 μg/ml

Host Species Mouse IgG1 Conjugate None

Endotoxin N.A. Purification Protein G

Storage 4°C

Application notes

	IHC-F	IHC-P	IF	FC	FS	IA	IP	W
Reference #								
Yes	•	•		•				
No							•	•
N.D.			•		•	•		

N.D.= Not Determined; IHC = Immuno histochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:10.

General Information

Description

The human anaphylatoxin C5a is a 74-residu glycopolypeptide which is generated by proteolytic cleavage of the complement factor C5 in the course of complement activation. A variety of biological effects evoked by C5a could be demonstrated, rendering this molecule an important mediator of inflammation, with granulocytes and macrophages as the main target cells. All cellular responses to C5a are specifically mediated by interactions with the membrane bound C5a receptor, a seven transmembrane GTP-binding-protein-coupled receptor that belongs to the rhodopsin supergene family. The C5a receptor is approximately 45 kDa.

References

- Opperman, M et al; Probing the human receptor for C5a anaphylatoxin with side-directed antibodies. J Immunol 1993. 151: 3785
- Opperman, M et al; Antibodies from the myeloid panel that react with the C5a receptor and antagonize C5a biological activity. Schlossman, S.F. (ed.) Leucocyte typing V.O.U.P.: 955-956
- Werfel, T et al; Binding of anti-C5a receptor (C5aR) antibodies to cells of clinically normal human skin. Schlossman, S.F. (ed.) Leucocyte typing V.O.U.P.: 957-959
- Opperman, M et al; Plasma clearance of human C5a anaphylatoxin by binding to leucocytes C5a receptors. Immunol 1994, 82: 516

Storage&stability

Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year.

Precautions

For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent infringements that might result from the use or derivation of this product.

We hereby certify that the above-stated information is correct and that this product has been successfully tested by the Quality Control Department. This product was released for sale according to the existing specifications. This document has been produced electronically and is valid without a signature.

Approved by Manager of QC Brenda Teunissen

Date 16/11/2020

Version: 08-2020

Do you have any questions or comments regarding this product? Please contact us via support@hycultbiotech.com.