

CERTIFICATE OF ANALYSIS – TECHNICAL DATA SHEET

Product name	MPO, Human, clone 266-6K1	Expiry date	-
Catalog number	HM2164-100UG	Amount	100 µg
Lot number	-	Concentration	100 µg/ml
Volume	1 ml	Conjugate	None
Formulation	0.2 µm filtered in PBS+0.1%BSA+0.02%NaN ₃	Purification	Protein G
Host Species	Mouse IgG1		
Endotoxin	N.A.		
Storage	4°C		

Application notes

	IHC-F	IHC-P	IF	FC	FS	IA	IP	W
Reference #								1
Yes						•		•
No		•						
N.D.	•		•	•	•		•	

N.D.= Not Determined; IHC = Immuno histochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.

- IA: the antibody can be used as coat and detection antibody.
- W: Samples were subjected to SDS-PAGE and transferred to PVDF membrane (Ref.1).
- Positive control: Neutrophils, HL-60 cells; Negative control: Erythrocytes.

General Information

Description	The monoclonal antibody 266-6K1 recognizes human myeloperoxidase (MPO), an ~135 glycoprotein expressed in all cells of the myeloid lineage. MPO functions as an α ₂ β ₂ heteromultimer consisting of two heavy (α) and two light (β) chains of 55 and 15 kDa respectively. MPO is abundantly present in azurophilic granules of polymorphonuclear neutrophils (PMNs). It is an important enzyme used during phagocytic lysis of engulfed foreign particles which takes part in the defense of the organism through production of hypochlorous acid (HOCl), a potent oxidant. In the stimulated PMN, MPO catalyzes the production of hypohalous acids, primarily hypochlorous acid in physiologic situations, and other toxic intermediates that greatly enhance PMN microbicidal activity. Upon activation of neutrophils, MPO can be rapidly released and as such useful in body fluids as marker for inflammatory status. Involvement of MPO has been described in numerous diseases such as atherosclerosis, lung cancer, Alzheimer's disease, inflammatory bowel disease and multiple sclerosis. Autoimmune antibodies to MPO (so called ANCA) are involved in Wegener's disease. Since the discovery of MPO deficiency, initially regarded as rare and restricted to patients suffering from severe infections, MPO has attracted more clinical attention.
Aliases	Myeloperoxidase
Cross reactivity	Rat: No; Mouse: No.
References	1. La Rocca, G et al; Oxidative stress induces myeloperoxidase expression in endocardial endothelial cells from patients with chronic heart failure. <i>Basic Res Cardiol</i> 2009, <i>104</i> : 307
Storage&stability	Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year.

Precautions

For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent infringements that might result from the use or derivation of this product.

We hereby certify that the above-stated information is correct and that this product has been successfully tested by the Quality Control Department. This product was released for sale according to the existing specifications. This document has been produced electronically and is valid without a signature.

Approved by Manager of QC
Brenda Teunissen

Date
03/12/2019

Do you have any questions or comments regarding this product? Please contact us via support@hycultbiotech.com.