

CERTIFICATE OF ANALYSIS – TECHNICAL DATA SHEET

Product name	Punctin-1, Human, clone 1B2-8-2		
Catalog number	HM2169-100UG		
Lot number	-	Expiry date	-
Volume	1 ml	Amount	100 µg
Formulation	0.2 µm filtered in PBS+0.1%BSA+0.02%NaN3	Concentration	100 µg/ml
Host Species	Mouse IgG2b	Conjugate	None
Endotoxin	N.A.	Purification	Protein G
Storage	4°C		

Application notes

	IHC-F	IHC-P	IF	FC	FS	IA	IP	W
Reference #								
Yes						•		•
No								
N.D.	•	•	•	•	•		•	

N.D.= Not Determined; IHC = Immuno histochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.

- W: For Western blotting applications the antibody works better in non-reducing than reducing conditions although it will also recognize the reduced antigen at correspondingly higher concentrations.

General Information

Description Monoclonal antibody 1B2-8-2 reacts with Punctin-1 (ADAMTSL-1). Punctin-1 is a secreted molecule resembling members of the family of proteases designated ADAMTS (a disintegrin-like and metalloprotease domain with thrombospondin type I motif). ADAMTS proteases are secreted enzymes that consist of a metalloprotease domain (which provides the catalytic activity) and ancillary domains (which provide substrate, cell or extracellular matrix activity). Punctin-1 lacks the pro-metalloprotease and the disintegrin-like domain typical of this family but contains other ADAMTS domains in precise order including 4 thrombospondin-I repeats. Punctin-1 is an extracellular matrix protein in tissues and deposited into the cell substratum and localized towards the periphery of transfected cells. Punctin-1 and Punctin-2 (ADAMTSL-3) are the first members of a novel family of ADAMTS-like proteins that appear to have important functions in the extracellular matrix. Monoclonal antibody 1B2-8-2 recognizes the first thrombospondin repeat near the N-terminal and reacts with molecular species of 55 kDa or a doublet of 55 kDa and 60 kDa (glycosylated form). Note that proteolysis of Punctin-1 has been described and additional smaller bands may also be obtained.

Aliases ADAMTSL-1

References 1. Hirohata, S et al; Punctin, a novel ADAMTS-like molecule, ADAMTSL-1, in extracellular matrix. J Biol Chem 2002, 277: 12182

Storage&stability Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year.

Precautions For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent infringements that might result from the use or derivation of this product.

We hereby certify that the above-stated information is correct and that this product has been successfully tested by the Quality Control Department. This product was released for sale according to the existing specifications. This document has been produced electronically and is valid without a signature.

Approved by Manager of QC
Brenda Teunissen

Date
03/12/2019

Do you have any questions or comments regarding this product? Please contact us via support@hycultbiotech.com.