

CERTIFICATE OF ANALYSIS – TECHNICAL DATA SHEET

Product name MPO, Human, clone B7

Catalog number HM2288-20UG

Lot number - Expiry date -

 $\begin{tabular}{lll} \begin{tabular}{lll} \begin$

Formulation 0.2 μm filtered in PBS+0.1%BSA+0.02%NaN3 Concentration 100 μg/ml

Host Species Mouse IgG1 Conjugate None


Endotoxin N.A. Purification Protein G

Storage 4°C

Application notes

	IHC-F	IHC-P	IF	FC	FS	IA	IP	W
Reference #								
Yes	•			•		•		•
No								
N.D.		•	•		•		•	

N.D.= Not Determined; IHC = Immuno histochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot



FC: Flow cytometric detection of MPO in erytrocytes (mAb B7, Cat.# HM2288). Red, green and blue line represents 1, 0.5 and 0.1 μ g/ml, respectively.

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.

- W: A reduced sample treatment and SDS-Page was used. The band size is 195 kDa (Ref FC: Antibody B7 stains MPO. For intracellular staining leukocytes were permeabilized with permeabilisation buffer of Invitrogen. The cells were fixed in formaldehyde before staining. As positive control LPS stimulated mouse erytrocytes was used and as negative control unstimulated mouse erythrocytes.
- W: A reduced sample treatment and SDS-Page was used. The band size is 63 kDa.
- IHC-F: Tissue sections were fixed in acetone and pretreated with PBS and 30 minutes with 2% BSA-PBS solution for blocking. As
 positive control spleen of a wild-type mouse was used and as negative control spleen tissue of a knock-out MPO mouse.
- Positive control: Neutrophils, HL-60 cells; Negative control: Cells stimulated with physiological saline.

General Information

Description

The monoclonal antibody B7 recognizes human myeloperoxidase (MPO), an ~135 glycoprotein expressed in all cells of the myeloid linage. MPO functions as an $\alpha2\beta2$ heteromultimer consisting of two heavy (α) and two light (β) chains of 55 and 15 kDa respectively. MPO is abundantly present in azurophilic granules of polymorphonuclear neutrophils (PMNs). It is an important enzyme used during phagocytic lysis of engulfed foreign particles which takes part in the defense of the organism through production of hypochlorous acid (HOCl), a potent oxidant. In the stimulated PMN, MPO catalyzes the production of hypohalous acids, primarily hypochlorous acid in physiologic situations, and other toxic intermediates that greatly enhance PMN microbicidal activity. Upon activation of neutrophils, MPO can be rapidly released and as such useful in body fluids as marker for inflammatory status. Involvement of MPO has been described in numerous diseases such as atherosclerosis, lung cancer, Alzheimer's disease, inflammatory bowel disease and multiple sclerosis. Autoimmune antibodies to MPO (so called ANCA) are involved in Wegener's disease. Since the discovery of MPO deficiency, initially regarded as rare and restricted to patients suffering from severe infections, MPO has attracted more clinical attention.

Version: 08-2020

Immunogen Mouse MPO.

Aliases Myeloperoxidase.

Cross reactivity Mouse: Yes; Rat: Yes.

Storage&stability Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year.

Precautions For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent

infringements that might result from the use or derivation of this product.

We hereby certify that the above-stated information is correct and that this product has been successfully tested by the Quality Control Department. This product was released for sale according to the existing specifications. This document has been produced electronically and is valid without a signature.

Approved by Manager of QC Brenda Teunissen

Date 09/12/2020

Do you have any questions or comments regarding this product? Please contact us via support@hycultbiotech.com.

Version: 08-2020