

CERTIFICATE OF ANALYSIS – TECHNICAL DATA SHEET

Product name	SP-B, Pig, pAb		
Catalog number	HP7002-20UG		
Lot number	-	Expiry date	-
Volume	200 µl	Amount	20 µg
Formulation	0.2 µm filtered in PBS+0.1%BSA+0.02%NaN3	Concentration	100 µg/ml
Host species	Rabbit IgG	Conjugate	None
Endotoxin	N.A.	Purification	Protein A
Storage	4°C		

Application notes

	IHC-F	IHC-P	IF	FC	FS	IA	IP	W
Reference #								
Yes						•		•
No								
N.D.	•	•	•	•	•		•	

N.D.= Not Determined; IHC = Immuno histochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.

- IA: The antibody can be used as capture and detection antibody.

General Information

Description	The polyclonal antibody recognizes the human surfactant protein B (SP-B). There are four surfactant-specific proteins, designated surfactant protein A (SP-A), SP-B, SP-C and SP-D respectively. SP-A and SP-D are hydrophilic surfactant proteins and are members of the collectin family. SP-B and SP-C are hydrophobic surfactant proteins and may be the most appropriate indicators for the evolutionary origin of surfactant. SP-B is synthesized by the alveolar type II epithelial cells as a 40-42 kD precursor that is subsequently proteolytically processed to 7.8-8 kD. SP-B enhances the spreading and stability of surfactant phospholipids in the alveolus. SP-B is essential for air-breathing in mammals and is therefore largely conserved. SP-B can interact with both phospholipid head groups and fatty chains and is particularly active in enhancing surface active behaviour in endogenous and exogenous lung surfactants. Even low SP-B contents had measurable effects in increasing the adsorption, dynamic surface tension lowering, and/or film respreading of DPPC, mixed synthetic lipids, and column-purified lung surfactant phospholipids. Deficiency of SP-B and other surfactant components is associated with respiratory distress syndrome (RDS) in premature infants and adults with respiratory distress syndrome (ARDS). The polyclonal antibody is raised against pig SP-B from pig lungs which has been reconstituted in micelles of lysophosphatidylcholine. The antibody recognizes under nonreducing conditions the most prominent forms of SP-B of 8 kD and 22 kD, corresponding to SP-B dimers. The polyclonal antibody is cross reactive with human SP-B.
Cross reactivity	Human: Yes.
References	<ol style="list-style-type: none"> Oviedo, J et al; Quantitation of pulmonary surfactant protein SP-B in the absence or presence of phospholipids by enzyme-linked immunosorbent assay. <i>Anal Biochem</i> 2001, 293: 78 Perez-Gil, J et al; Lipid-protein interactions of hydrophobic proteins SP-B and SP-C in lung surfactant assembly and dynamics. <i>Pediatr Pathol Mol Med</i> 2001, 20: 445
Storage&stability	Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year.

Precautions

For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent infringements that might result from the use or derivation of this product.

We hereby certify that the above-stated information is correct and that this product has been successfully tested by the Quality Control Department. This product was released for sale according to the existing specifications. This document has been produced electronically and is valid without a signature.

Approved by Manager of QC
Brenda Teunissen

Date
13/01/2021

Do you have any questions or comments regarding this product? Please contact us via support@hycultbiotech.com.