

CERTIFICATE OF ANALYSIS – TECHNICAL DATA SHEET

Product name	TLR2, Mouse, pAb	Expiry date	-
Catalog number	HP8016-20UG	Amount	20 µg
Lot number	-	Concentration	100 µg/ml
Volume	200 µl	Conjugate	None
Formulation	0.2 µm filtered in PBS+0.1%BSA+0.02%NaN3	Purification	Affinity
Host species	Rabbit Ig		
Endotoxin	N.A.		
Storage	4°C		

Application notes

	IHC-F	IHC-P	IF	FC	FS	IA	IP	W
Reference #								
Yes				•				•
No								
N.D.	•	•	•		•	•	•	

N.D.= Not Determined; IHC = Immuno histochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.

General Information

Description	The polyclonal antibody reacts with mouse Toll-like receptor 2 (TLR2 (CD282)). Toll-like receptors (TLR) are highly conserved throughout evolution and have been implicated in the innate defense to many pathogens. In <i>Drosophila</i> toll is required for the anti-fungal response, while the related 18-wheeler is involved in antibacterial defenses. In mammals, TLR identified as type I transmembrane signaling receptors with pattern recognition capabilities, have been implicated in the innate host defense to pathogens. TLR2 has been identified as a receptor that is central to the innate immune response to lipoproteins of Gram-negative bacteria, several whole Gram-positive bacteria, as well as a receptor for peptidoglycan and lipoteichoic acid and other bacterial cell membrane products. A functional interaction between TLR2 and TLR6 in the cellular response to various bacterial products has been discovered. The currently accepted paradigm regards TLR2 as an essential receptor for many eubacterial cell wall components, including lipoproteins and peptidoglycan. Bacterial species as diverse as mycobacteria, spirochetes, mycoplasma, <i>Staphylococcus aureus</i> , and <i>Streptococcus pneumoniae</i> have all been shown to mediate cellular activation via TLR2.
References	1. Sing, A et al; <i>Yersinia</i> V-antigen exploits toll-like receptor 2 and CD14 for interleukin 10-mediated immunosuppression. <i>J Exp Med</i> 2002, 21: 1017
Storage&stability	Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year.
Precautions	For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent infringements that might result from the use or derivation of this product.

We hereby certify that the above-stated information is correct and that this product has been successfully tested by the Quality Control Department. This product was released for sale according to the existing specifications. This document has been produced electronically and is valid without a signature.

Approved by Manager of QC
Brenda Teunissen

Date
13/01/2021

Do you have any questions or comments regarding this product? Please contact us via support@hycultbiotech.com.