

## **CERTIFICATE OF ANALYSIS – TECHNICAL DATA SHEET**

Catalog numberHP8017-20UGLot number-Colume20 μl20 μlAmountPormulation0.2 μm filtered in PBS+0.1%BSA+0.02%NaN3Most speciesRabbit IgGConjugateNoneEndotoxinN.A.PurificationProtein A	Product name	VG5Q, Mouse, pAb		
Volume200 μlAmount20 μgFormulation0.2 μm filtered in PBS+0.1%BSA+0.02%NaN3Concentration100 μg/mlHost speciesRabbit IgGConjugateNoneEndotoxinN.A.PurificationProtein A	Catalog number	HP8017-20UG		
Formulation0.2 μm filtered in PBS+0.1%BSA+0.02%NaN3Concentration100 μg/mlHost speciesRabbit IgGConjugateNoneEndotoxinN.A.PurificationProtein A	Lot number	-	Expiry date	-
Host speciesRabbit IgGConjugateNoneEndotoxinN.A.PurificationProtein A	Volume	200 μΙ	Amount	20 µg
EndotoxinN.A.PurificationProtein A	Formulation	0.2 μm filtered in PBS+0.1%BSA+0.02%NaN3	Concentration	100 µg/ml
	Host species	Rabbit IgG	Conjugate	None
Storage 4°C	Endotoxin	N.A.	Purification	Protein A
	Storage	4°C		

## **Application notes**

	IHC-F	IHC-P	IF	FC	FS	IA	IP	W
Reference #								
Yes	•	•				•	•	•
No								

N.D.= Not Determined; IHC = Immuno histochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:10.

## **General Information**

Description	The polyclonal antibody reacts with mouse VG5Q, a 84 kDa protein. VG5Q functions as an angiogenic factor i promoting angiogenesis and suppression of VG5Q expression inhibits vessel formation. Angiogenic factors are critica to the initiation of angiogenesis and maintenance of the vascular network. Angiogenesis has an essential role i pathological conditions such as cancer and various ischaemic and inflammatory diseases. VG5Q can bind t endothelial cells and promote cell proliferation, suggesting that the protein may act in an autocrine fashion. VG5G interacts with TWEAK (also known as TNFSF12), another secreted angiogenic factor. VG5Q shows strong expressio in blood vessels and is secreted when vessel formation is initiated. VG5Q protein was detected mostly in the cytoplasr and around the nuclei of human microvascular endothelial cells (HMVECs). Furthermore VG5Q is detected in huma umbilical vein endothelial cells (HUVECs), human heart fibroblast (HHF) and ovarian cancer cells (OV-3), but low expression was detected in kidney cancer cells (RP-45), HeLa Cells and bladder cancer cells.			
References	1. Tian, X et al; Identification of an angiogenic factor that when mutated causes susceptibility to Klippel-Trenaunay syndrome. Nature 2004, <i>427</i> : 640			
Storage&stability	Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year.			
Dressutions	For research use only. Not for use in or on humans or enimple or for discretion. It is the representibility of the user to			

Precautions For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent infringements that might result from the use or derivation of this product.

We hereby certify that the above-stated information is correct and that this product has been successfully tested by the Quality Control Department. This product was released for sale according to the existing specifications. This document has been produced electronically and is valid without a signature.

Approved by Manager of QC Brenda Teunissen

Date 13/01/2021

Do you have any questions or comments regarding this product? Please contact us via support@hycultbiotech.com.