

CERTIFICATE OF ANALYSIS – TECHNICAL DATA SHEET

Product name	VG5Q, Human, pAb		
Catalog number	HP9037-20UG		
Lot number	-	Expiry date	-
Volume	200 μΙ	Amount	20 µg
Formulation	0.2 μm filtered in PBS+0.1%BSA+0.02%NaN3	Concentration	100 μg/ml
Host Species	Rabbit IgG	Conjugate	None
Endotoxin	N.A.	Purification	Protein A
Storage	4°C		

Application notes

	IHC-F	IHC-P	IF	FC	FS	IA	IP	W
Reference #								
Yes	•	•				•	•	٠
No								
N.D.			•	•	•			

N.D.= Not Determined; IHC = Immuno histochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.

General Information			
Description	The polyclonal antibody reacts with human VG5Q, a 84 kDa protein. VG5Q functions as an angiogenic factor promoting angiogenesis and suppression of VG5Q expression inhibits vessel formation. Angiogenic factors are crit to the initiation of angiogenesis and maintenance of the vascular network. Angiogenesis has an essential role pathological conditions such as cancer and various ischaemic and inflammatory diseases. VG5Q can bind endothelial cells and promote cell proliferation, suggesting that the protein may act in an autocrine fashion. VG interacts with TWEAK (also known as TNFSF12), another secreted angiogenic factor. VG5Q shows strong express in blood vessels and is secreted when vessel formation is initiated. VG5Q protein was detected mostly in the cytopla and around the nuclei of human microvascular endothelial cells (HMVECs). Furthermore VG5Q is detected in hur umbilical vein endothelial cells (HUVECs), human heart fibroblast (HHF) and ovarian cancer cells (OV-3), but expression was detected in kidney cancer cells (RP-45), HeLa Cells and bladder cancer cells		
References	1. Tian, X et al; Identification of an angiogenic factor that when mutated causes susceptibility to Klippel-Trenau syndrome. Nature 2004, <i>427</i> : 640		
Storage&stability	Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year.		
Precautions	For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the use comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any pa infringements that might result from the use or derivation of this product.		

Approved by Manager of QC Brenda Teunissen

Date 31/03/2021

Do you have any questions or comments regarding this product? Please contact us via support@hycultbiotech.com.