


CERTIFICATE OF ANALYSIS – TECHNICAL DATA SHEET

Product name	Beta-defensin 2, Human, pAb		
Catalog number	HP9057-20UG		
Lot number	-	Expiry date	-
Volume	200 µl	Amount	20 µg
Formulation	0.2 µm filtered in PBS+0.1%BSA+0.02%NaN3	Concentration	100 µg/ml
Host Species	Rabbit Ig	Conjugate	None
Endotoxin	N.A.	Purification	Protein A
Storage	4°C		

Application notes

	IHC-F	IHC-P	IF	FC	FS	IA	IP	W
Reference #								
Yes		•						•
No						•		
N.D.	•		•	•	•		•	

N.D.= Not Determined; IHC = Immuno histochemistry; F = Frozen sections; P = Paraffin sections; IF = Immuno Fluorescence; FC = Flow Cytometry; FS = Functional Studies; IA = Immuno Assays; IP = Immuno Precipitation; W = Western blot



IHC-P: Immunohistochemical analysis of beta-defensin 2 in paraffin-embedded human lung tissue using pAb HP9057.

Dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.

- W: A reduced sample treatment and SDS-Page was used. The band size is 25kDa for recombinant protein with tag and 5 kDa for the native protein.
- IHC-P: Tissue sections were pretreated with high temperature citrate. Tissue sections were fixed in formalin. As a negative control PBS was used.

General Information

Description	<p>The polyclonal antibody recognizes human beta-defensin 2 (hBD-2). hBD-2 is a cystein-rich cationic 41 amino acid antimicrobial peptide of 4-5 kDa. Human BD-2 is produced by epithelial cells upon stimulus by lipopolysaccharides and proinflammatory cytokines TNFα and IL1β. Contact of keratinocytes with gram-negative bacteria results in rapid induction of hBD-2 protein.</p> <p>hBD-2 has been described as a dynamic component of the local epithelial defense system of the skin, intestinal and respiratory tract, where it functions by protecting surfaces from infection. Its local expression has been associated with skin lesions like psoriasis as well as infected lung epithelia of patients with cystic fibrosis. Furthermore, in inflammatory bowel disease (IBD), expression of hBD-2 is increased in patients with IBD compared to healthy persons.</p> <p>The NF-kB pathway has been recognized as a key component in the induction of hBD-2 expression, however, other studies have observed induction mediated by the mitogen-activated protein kinase (MAPK) pathways. Thus, increased expression of hBD-2 in epithelial cells is associated with the proinflammatory response. This is supported by the finding that the anti-inflammatory cytokines IL-10 and IL-13 downregulate the synthesis of hBD-2 in atopic dermatitis.</p>
Immunogen	CHPVFCPRRYKQIGTCGL
Aliases	beta-defensin 2, hBD-2

Storage&stability Product should be stored at 4°C. Under recommended storage conditions, product is stable for at least one year.

Precautions For research use only. Not for use in or on humans or animals or for diagnostics. It is the responsibility of the user to comply with all local/state and federal rules in the use of this product. Hycult Biotech is not responsible for any patent infringements that might result from the use or derivation of this product.

We hereby certify that the above-stated information is correct and that this product has been successfully tested by the Quality Control Department. This product was released for sale according to the existing specifications. This document has been produced electronically and is valid without a signature.

Approved by Manager of QC
Brenda Teunissen

Date
31/03/2021

Do you have any questions or comments regarding this product? Please contact us via support@hycultbiotech.com.